

**Porkert et al. - Timing of breeding in Common Redstart populations**

**Table A4.** Means of Spearman correlation coefficients of the daily number of started broods and the mean air temperature over the previous 3, 5 and 10 days in each year in the studied populations (Rt3, Rt5 and Rt10, resp.). ID numbers of populations as in Table 1.

ID	Country	<i>n</i>	Rt3	<i>SE</i>	Rt5	<i>SE</i>	Rt10	<i>SE</i>
1	GER	14	-0.1334	0.0883	-0.0427	0.0852	-0.0331	0.1137
2	CZE	14	0.0821	0.0918	0.1776	0.1040	0.0245	0.0927
3	RUS	5	0.0929	0.2042	0.1324	0.1779	0.0949	0.1893
4	RUS	19	0.0254	0.0708	-0.0314	0.0858	-0.0664	0.0634
5	RUS	17	0.2134	0.1040	0.1217	0.1113	0.0810	0.0835
6	FIN	24	0.1939	0.0799	0.2354	0.0739	-0.0355	0.0679
7	FIN	23	0.0031	0.0594	0.0554	0.0664	-0.1089	0.0603
8	FIN	9	0.2122	0.1264	0.1566	0.1377	0.0895	0.1398
9	FIN	32	0.0753	0.0610	0.0677	0.0617	-0.0054	0.0647