

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/328759720>

Ecology and Geography of Plants and Plant Communities The fourth International Scientific Conference on Ecology and Geography of Plants and Plant Communities Volume 2018 Lichen-lik...

Conference Paper · October 2018

DOI: 10.18502/kls.v4i7.3231

CITATIONS

0

READS

21

4 authors:



Victor Andreevich Mukhin

Russian Academy of Sciences

88 PUBLICATIONS 193 CITATIONS

[SEE PROFILE](#)



Nadezhda Neustroeva

Russian Academy of Sciences

8 PUBLICATIONS 4 CITATIONS

[SEE PROFILE](#)



Elena Patova

Komi Scientific Center

49 PUBLICATIONS 173 CITATIONS

[SEE PROFILE](#)



Irina V. Novakovskaya

Komi Scientific Center

17 PUBLICATIONS 10 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Wood plants water, carbon and nitrogen status under climate change [View project](#)



Charophytes (Charophyceae, Charales) of Eastern Europe, Western, Northern and Central Asia [View project](#)

Conference Paper

Lichen-like Symbiotic Associations of Wood-decaying Fungi and Algae. I. Biodiversity and Ecology of Photobionts

V. A. Mukhin^{1,2}, N. V. Neustroeva², E. N. Patova³, and I. V. Novakovskaya³¹Institute of Natural Sciences, Ural Federal University, 620000 Ekaterinburg, Russia²Institute of Plant and Animal Ecology, Ural Branch of the Russian Academy of Sciences, 620144 Ekaterinburg, Russia³Institute of Biology, Komi Scientific Center, Ural Branch of the Russian Academy of Sciences, 167982 Syktyvkar, Russia

Abstract

The article presents new data on the taxonomical, morphological and ecological composition and species diversity of symbiont algae associated with xylotrophic fungi. The largest part of symbionts (86%) are eukaryotic algae belonging to the divisions *Chlorophyta* (68% of total number of species), *Ochrophyta* (9%) and *Charophyta* (8%). The prokaryotic algae, or *Cyanoprokaryota*, make up the remaining 14% of species. The eukaryotic algae are an obligatory component of mycetobiont communities, whereas *Cyanoprokaryota* are the optional, facultative part. Out of 46 mycetobiont algae genera, 29 (or 64%) are single-species taxa, while 15 (32%) genera include two or three species. Two genera – *Chlamydomonas* and *Klebsormidium* – are represented by 6 and 4 species, respectively. The majority of mycetobiont algae have coccoid (41%) and trichal (33%) thalli, colonial-coccoid (18%) and monadic (8%) algae are rarely observed. All algae species belong to widespread epiphytic, soil and lichenophilic groups that do not require symbiosis with fungi. Obligatory mycetobionts were not observed during the study. Communities of mycetobiont algae have host-specificity and high geographical and individual variability.

Keywords: wood-decaying fungi, algae and *Cyanoprokaryota*, biodiversity, ecology, symbiosis

1. Introduction

The widely observed presence of algae in basidiocarps of xylotrophic basidiomycetes remains poorly studied. Only a few articles address the issue. In one of the first related studies, Burdsall and Volk [1] reported two species of one-celled algae observed in basidiocarps of *Oxyporus nobilissimus* W. B. Cooke. In articles by Zavada, Simoes [2]

Corresponding Author:

V. A. Mukhin
victor.mukhin@ipae.uran.ru

Received: 12 September 2018

Accepted: 15 October 2018

Published: 29 October 2018

Publishing services provided by
Knowledge E

© V. A. Mukhin et al. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Selection and Peer-review under the responsibility of the Ecology and Geography of Plants and Plant Communities Conference Committee.

 OPEN ACCESS

and Zavada et al. [3], 4 species of green algae (*Hormidium* sp., *Stichococcus bacillaris* Nägeli, *Chlorococcum* sp. and *Trebouxia* sp.) were found in basidiocarps of *Trametes versicolor* (L.) Lloyd. In another work by Videv et al. [4] reported 10 species: 9 belonged to green algae and one, *Klebsormidium dissectum* (F. Gray) Y. Ettl. et Gärtner, to Charophyta. Stoyneva et al. [5] described the growth of 4 green algae species, *Desmococcus vulgaris*, *Trebouxia arboricola*, *Stichococcus bacillaris* and *S. minutus*, in basidiocarps of *Fomes fomentarius* (L.) Fr. In total, 16 taxons of algae are found in basidiocarps of xylotrophic fungi, according to mentioned articles. Their interaction type with the fungi remains under a question. Some researchers [4, 5] say that algae developing on basidiocarps cannot be viewed as symbionts but should be rather seen as epiphytic, epimicotic organisms. However, there is convincing evidence of symbiosis between algae and *T. versicolor*, referred to as 'possible demi-lichenization of the basidiocarps' [2, 3], an interaction which is similar to those that mycobionts and photobionts have in lichens. We think that the relationship between mycetobiont algae and fungi should be characterized as 'associative symbiosis', in which the algae receives protection from excessive insolation and CO₂ and H₂O from respiration and, in return, the fungus gets an additional source of carbon and nitrogen nutrition to wooden substrate [6]. In Zavada et al.'s [3] article and our own works, it has been experimentally proven that products of algal photosynthesis are found in fungi [7, 8] and that the fixation of molecular nitrogen happens when prokaryotic blue-green algae (Cyanoprokaryota) are present in basidiocarps [9].

Therefore, associations between wood-destroying fungi and algae require the most careful study, as this symbiosis has unique and significant importance for a biosphere. The biodiversity and ecology of symbiotic algae in such associations should be given special attention.

2. Methods

The objects of research mycetobiont algae in basidiocarps of wood-destroying fungi, were collected from four locations: (a) pine-birch pre-forest-steppe (56°36'5" N, 61°3'24" E), (b) dark coniferous forests of southern taiga, the Central Urals (57°21'01" N, 58°41'54" E), (c) the Valdai Upland (57°57.76' N, 33°20.34' E) and (d) taiga forests of the Komi Republic (61°34' N, 50°36' E). Before microscopic examination of the algae, the basidiocarps were moistened with distillate for two days. After we divided them into two parts, each got a different treatment for 7 days: one was moistened with 3N BBM nutrient medium for green algae, while a Bg11 nutrient medium for *Cyanoprokaryota*

was applied to the other. The algae (surface abrading probes and sections of a basidiocarp) were examined under a Nikon Eclipse 80i microscope (magnification x400, x1000). For better identification of algal species diversity, we transferred fragments of basidiocarps and washings from their surface into agar and liquid nutrient media cultures: 3N BBM and Bg11 were used for green algae and *Cyanoprokaryota*, respectively. Algae were identified using Russian and foreign published literature and taxonomic keys [10–12]. The part of the species which requires observations during different stages of reproduction and development were determined only to the genus level.

3. Results

Our findings (Table 1) show that mycetobiont algae are found in the basidiocarps of many xylotrophic basidiomycetes, though they mostly inhabit annual or overwintering basidiocarps: very rarely, they inhabit perennial ones. Fungi with algae in basidiocarps have a diverse ecology: they decompose deciduous and coniferous wood, cause white and brown rot. The *Cerrena unicolor* and *Trichaptum pargamenum* species have the highest diversity of mycetobiont algae (31–34 species). The basidiocarps of *Stereum subtomentosum*, *Trichaptum abietinum* and *T. fuscoviolaceum* have 20–22 species of algae. 11 to 15 symbiont algae species are found in *Stereum hirsutum*, *Trametes gibbosa*, *T. ochracea* and *T. pubescens*. Most fungi have no more than 10 species of algae in their basidiocarps. The data analysis shows that eukaryotic algae are obligatory components in basidiocarps of all fungi species, whereas *Cyanoprokaryota* serve as a small and additional (facultative) part. Eukaryotic algae mainly or exclusively belong to the *Chlorophyta* division in most fungi species: only in one, *Gloeophyllum sepiarium*, are members from the group absent (Table 1). As shown earlier by our research [6], the proportion of basidiocarps with algae varies from 30% to 90% between fungi species, indicating that such a relationship is not obligatory.

The taxonomical and morphological characteristics of mycetobiont algae are presented in Table 2. The largest number of species (64% or 86%) belong to eukaryotic algae from the *Chlorophyta* (51 species, 68%), *Ochrophyta* (7 species, 9%) and *Charophyta* (6 species, 8%) divisions: 15% of species belong to *Cyanoprokaryota*. From 46 genera of mycetobiont algae, 29 (or 64%) consist of one species, while 15 have two or three species. *Chlamydomonas* and *Klebsormidium* are the most diverse genus, with 6 and 4 species, respectively. The average species diversity within genus is 1.7.

TABLE 1: The list of xylotrophic fungi with mycetobiont algae in basidiocarps (total number of species/number of species belonging to *Chlorophyta/Charophyta/Ochrophyta/Cyanoprokaryota* divisions, respectively).

Xylotrophic Fungi	Mycetobiont Algae
<i>Bjerkandera adusta</i> (Willd.) P. Karst. (awd)*	9 / 5 / 0 / 1 / 3
<i>Cerrena unicolor</i> (Bull.) Murrill (awd)	31 / 23 / 1 / 4 / 3
<i>Chondrostereum purpureum</i> (Pers.) Pouzar (awd)	7 / 7 / 0 / 0 / 0
<i>Datronia mollis</i> (Sommerf.) Donk (awd)	3 / 3 / 0 / 0 / 0
<i>Echlerialla deglubens</i> (Berk. & Br.) D.A. Reid (awd)	3 / 3 / 0 / 0 / 0
<i>Fomes fomentarius</i> (L.) Fr. (pwd)	2 / 2 / 0 / 0 / 0
<i>Gloeophyllum sepiarium</i> (Wulfen) P. Karst. (abc)	4 / 0 / 1 / 1 / 2
<i>Lenzites betulina</i> (L.) Fr. (awd)	8 / 7 / 0 / 0 / 1
<i>Onnia leporina</i> (Fr.) H. Jahn (awc)	4 / 3 / 0 / 0 / 1
<i>Phellinus chrysoloma</i> (Fr.) Donk (pwc)	4 / 3 / 0 / 0 / 1
<i>Ph. ignarius</i> (L.) Quél. (pwd)	2 / 2 / 0 / 0 / 0
<i>Ph. tremulae</i> (Bondartsev) Bondartsev & P. N. Borisov (pwd)	3 / 2 / 0 / 0 / 1
<i>Steccherinum ochraceum</i> (Pers.) Gray (awd)	7 / 7 / 0 / 0 / 0
<i>Stereum hirsutum</i> (Willd.) Pers. (awd)	11 / 10 / 1 / 0 / 0
<i>S. subtomentosum</i> Pouzar (awd)	21 / 17 / 2 / 1 / 1
<i>Trametes gibbosa</i> (Pers.) Fr. (awd)	13 / 12 / 0 / 0 / 1
<i>T. hirsuta</i> (Wulfen) Lloyd (awd)	7 / 7 / 0 / 0 / 0
<i>T. ochracea</i> (Pers.) Gilb. & Ryvarden (awd)	15 / 10 / 0 / 1 / 4
<i>T. pubescens</i> (Schumach.) Pilát (awd)	12 / 8 / 1 / 1 / 1
<i>T. trogii</i> Berk. (awd)	9 / 6 / 1 / 2 / 0
<i>T. versicolor</i> (L.) Lloyd (awd)	9 / 7 / 1 / 1 / 0
<i>Trichaptum abietinum</i> (Dicks.) Ryvarden (awc)	22 / 18 / 2 / 2 / 0
<i>T. fuscoviolaceum</i> (Ehrenb.) Ryvarden (awc)	20 / 18 / 1 / 0 / 1
<i>T. pargamenum</i> (Fr.) G. Cunn. (awd)	34 / 28 / 4 / 2 / 0
Source: Authors' own work.	
Note: * – annual basidiocarp (a), perennial (p); brown rot (b), white rot (w); decomposers of deciduous (d) or coniferous (c) wood.	

Most algae have a coccoid (41%) and trichal (33%) thallus: colonial-coccoid (18%) and monadic (8%) forms are rare. All mycetobiont species are eurybiotic with a wide distribution: pedobionts and hydrobionts (*Chlamydomonas sp.*, etc.), epiphytes (*Desmococcus olivaceus*, etc.), aerophytes and photobionts of ascomycete lichens (*Pseudococcomyxa simplex*, *Stichococcus bacillaris*, etc.). There are no obligatory mycetobiont algae, which indicates the facultative nature of the relationship between the algae and fungi.

Only a few mycetobiont algae have a low degree of selectivity and can be found in many fungi species. For example, the *Interfilum terricola*, *Pseudococcomyxa simplex* and *Stichococcus bacillaris* species occur in basidiocarps of all fungi from the *Trametes*

TABLE 2: Taxonomical and morphological characteristics of mycetobiont algae dwelling in basidiocarps of xylotrophic fungi.

Genus	Number Species, thallus type *	Genus	Number Species, thallus type
<i>Chlorophyta</i>		<i>Chlorophyta</i>	
<i>Bracteacoccus</i>	1, c	<i>Tetracystis</i>	2, cc
<i>Chlamydomonas</i>	6, m	<i>Trebouxia</i>	2, c
<i>Chlorella</i>	3, c	<i>Ulothrix</i>	2, t
<i>Chlorococcum</i>	1, c	<i>Ochrophyta</i>	
<i>Chloroidium</i>	2, c	<i>Bumilleriopsis</i>	1, t
<i>Chlorosarcinopsis</i>	1, cc	<i>Characiopsis</i>	2, c
<i>Coenochloris</i>	3, cc	<i>Eustigmatos</i>	1, c
<i>Coenocystis</i>	1, cc	<i>Heterococcus</i>	1, t
<i>Desmococcus</i>	1, cc	<i>Tribonema</i>	1, t
<i>Dictyochloropsis</i>	1, c	<i>Vischeria</i>	1, c
<i>Diplosphaera</i>	1, cc	<i>Charophyta</i>	
<i>Elliptochloris</i>	3, c	<i>Cylindrocystis</i>	1, c
<i>Interfilum</i>	2, t	<i>Klebsormidium</i>	4, t
<i>Leptosira</i>	2, t	<i>Mesotaenium</i>	1, c
<i>Mychonastes</i>	1, c	<i>Cyanoprokaryota</i>	
<i>Myrmecia</i>	3, c	<i>Anabaena</i>	1, t
<i>Neochlorosarcina</i>	1, cc	<i>Aphanocapsa</i>	1, cc
<i>Neocystis</i>	1, cc	<i>Calothrix</i>	1, t
<i>Neospongiococcum</i>	3, c	<i>Chroococcus</i>	1, c
<i>Parietochloris</i>	1, c	<i>Desmonostoc</i>	1, t
<i>Pseudococcomyxa</i>	2, c	<i>Hassallia</i>	1, t
<i>Scotiellopsis</i>	1, c	<i>Nostoc</i>	3, t
<i>Spongiococcum</i>	1, cc	<i>Phormidium</i>	1, t
<i>Sporotetras</i>	1, cc	<i>Scytonema</i>	1, t
<i>Stichococcus</i>	2, t		

Note: * - coccoid (c), colonial-coccoid (cc), monadic (m), trichal (t).

genus (Table 3). However, the opposite group of algae, found only in 1 or 2 fungi species from a genus, is also observed. The members of this group are the green algae species *Diplosphaera chodatii*, *S. minor*, *Cyanoprokaryota* – *Hassallia byssoidea*, *Nostoc commune* and *Scytonema ocellatum*. Most likely, this group consists of both random and, possibly, highly selective species. In any case, they determine the specificity of mycetobiont algae communities [13]. This can be seen in the variation of mycetobiont algae richness between fungi species, particularly in the genus *Trametes* where the number of associated algae species varies from 7 to 15 species (Table 3).

The species composition of mycetobiont algae varies between basidiocarps belonging to the same fungi species: in other words, high individual variability is present. For example, basidiocarps of *Trichaptum pargamenum* (collected at the same time and in the same place, the Komi Republic, the middle-taiga forests) could have from 5 to 10 species of algae, but only 2 – *Pseudococcomyxa simplex* and *Stichococcus bacillaris* – are present in all basidiocarps. Such species as *Dictyochloropsis* sp., *Interfilum terricola*, *Klebsormidium pseudostichococcus*, *Leptosira* sp., *Trebouxia* sp. and *Vischeria helvetica*, are noted only in one of the basidiocarps. The majority of species (*Chlamydomonas* sp., *Coenocystis oleifera* (Broady) Hindák, *Desmococcus olivaceus*, *Diplosphaera chodatii*, *Elliptochloris* sp., *E. subsphaerica*, *Klebsormidium nitens* (Kützing) Lokhorst, *Myrmecia bisecta* Reisigl and *Sporotetras polydermatica*) occur in two or three basidiocarps of *Trichaptum pargamenum*. Therefore, each basidiocarp has its own particular algal composition – the Sørensen-Chekanovsky species similarity coefficient lies at the range from 0.3 to 0.6.

The geographic variability of mycetobiont algae composition is also observed [14]. For example, out of 16 species of algae found in the *Trichaptum pargamenum* basidiocarps from the middle taiga forests of the Komi Republic, only 6 occur in the basidiocarps of the same species in the Central Urals: *Chlamydomonas* sp., *Interfilum terricola*, *Desmococcus olivaceus*, *Pseudococcomyxa simplex*, *Stichococcus bacillaris* and *Trebouxia* sp. High individual and geographical variability, again, proves the non-obligatory connection of algae and fungus.

TABLE 3: Mycetobiont algae of the basidiocarps of the *Trametes* genus species: 1 – *T. gibbosa*, 2 – *T. hirsuta*, 3 – *T. ochracea*, 4 – *T. pubescens*, 5 – *T. trogii*, 6 – *T. versicolor*.

Mycetobiont algae	Species					
	1	2	3	4	5	6
<i>Chlorophyta</i>						
<i>Chlamydomonas</i> sp.			+	+	+	
<i>Chlorella vulgaris</i> f. <i>globosa</i> V. M. Andreyeva	+					
<i>Chloroidium saccharophilum</i> (W. Krüger) Darienko, Gustavs, Mudimu, Menendez, Schumann, Karsten, Friedl & Proschold	+	+	+			
<i>Coenochloris oleifera</i> (Broady) I. Kostikov, T. Darienko, A. Lukesová, & L. Hoffmann	+					
<i>Desmococcus olivaceus</i> (Persoon ex Acharius) J. R. Laundon	+	+		+		
<i>Diplosphaera chodatii</i> Bialosukniá	+				+	+
<i>Elliptochloris reniformis</i> (S. Watanabe) H. Ettl & G. Gärtner				+		+
<i>E. subsphaerica</i> (Reisigl) Ettl & Gärtner	+					

Mycetobiont algae	Species					
<i>Interfilum terricola</i> (J. B. Petersen) Mikhailyuk, Sluiman, Massalski, Mudimu, Demchenko, Friedl & Kondratyuk	+	+	+	+	+	+
<i>Leptosira</i> sp.			+		+	
<i>Mychonastes homosphaera</i> (Skuja) Kalina et Puncochárová					+	
<i>Myrmecia incisa</i> Reisingl					+	
<i>Myrmecia</i> sp.						+
<i>Neochlorosarcina</i> sp.	+					
<i>Pseudococcomyxa simplex</i> (Mainx) Fott	+	+	+	+	+	+
<i>Sporotetras polydermatica</i> (Kützing) I. Kostikov, T. Darienko, A. Lukesová, & L. Hoffmann			+	+	+	
<i>Stichococcus bacillaris</i> Nägeli	+	+	+	+	+	+
<i>S. minor</i> Nägeli	+		+			
<i>Tetracystis macrostigmata</i> Nakano					+	
<i>Trebouxia</i> sp.	+					+
<i>Ulothrix variabilis</i> Kützing					+	
Ochrophyta						
<i>Bumilleriopsis</i> sp.					+	
<i>Characiopsis</i> sp.				+		+
<i>Eustigmatos magnus</i> (J. B. Petersen) D. J. Hibberd						+
<i>Vischeria helvetica</i> (Vischer & Pascher) D. J. Hibberd						+
Charophyta						
<i>Klebsormidium pseudostichococcus</i> (Heering) H. Ettl & Gärtner					+	+
<i>Mesotaenium</i> sp.						+
Cyanoprokaryota						
<i>Desmonostoc muscorum</i> (C. Agardh ex Bornet & Flahault) Hrouzek & Ventura	+					
<i>Hassallia byssoidea</i> Hassal ex Bornet et Flahault					+	
<i>Nostoc</i> cf. <i>punctiforme</i> (Kützing ex Hariot) Hariot					+	
<i>N. commune</i> Vaucher ex Bornet et Flahault					+	
<i>Scytonema ocellatum</i> Lyngbye ex Bornet & Flahault					+	+
The number of species of algae in basidiocarps	13	7	15	12	9	9
Source: Authors' own work.						

4. Conclusion

Xylotrophic *Basidiomycetes* and mycetobiont algae inhabiting of their basidiocarps form multicomponent symbiotic associations consisting of a fungus (mycobiont) and several species of algae associated with it (photobionts). Mycobionts, as a rule, are fungi

with an annual, annual-wintering basidiocarps, and less often – perennial. The main, obligatory component of photobionts is green algae: other algal groups, such as *Ochrophyta*, *Charophyta* and *Cyanoprokaryota* are optional, not obligatory. The composition of mycetobiont algae in basidiocarps is not stable. It varies between different fungi species, the regions where they grow and within individuals. Although the symbiotic relationship between fungi and mycetobiont algae is not obligatory, it is beneficial for both: the algae have some protection from the environment and receive additional H₂O and CO₂ during their hosts' respiration, while the fungi get an additional source of carbon and nitrogen nutrition. In our opinion, the symbiotic associations of xylotrophic fungi and mycetobiont algae correspond to our notions of basidial lichens and should be considered as such.

Funding

The work has been conducted under the federal government assignment for the Institute of Plant and Animal Ecology UB RAS and partly with the support of the grant from Government of the Russian Federation (Act 211, Agreement 02.A03.21.0006) and RFBR grant № 18-04-00643.

References

- [1] Burdsall, H. H., Volk, T. J., and J. F. (1996). Ammirati, *Bridgeoporus*, a new genus to accommodate *Oxyporus nobilissimus* (Basidiomycota, Polyporaceae). *Mycotaxon*, vol. 60, pp. 387–395.
- [2] Zavada, M. S. and Simoes, P. (2001). The possible demi-lichenization of the basidiocarps of *Trametes versicolor* (L.: Fries) Pilat (Polyporaceae), *Northeast. Nature*, vol. 8, no. 1, pp. 101–112.
- [3] Zavada, M. S., DiMichele, L., and Toth, C. R. (2004). The possible demi-lichenization of *Trametes versicolor* (L.: Fr.) Pilat (Polyporaceae): The Transfer of Fixed ¹⁴CO₂ from Epiphytic Algae to *T. versicolor*, *Northeast. Nature*, vol. 11, no. 1, pp. 33–40.
- [4] Videv, P. V., Gärtner, G., Uzunov, B. A., et al. (2017). Epimycotic algae on the medicinal fungus *Trametes versicolor* (L.) Lloyd. *International Journal of Advanced Research in Botany (IJARB)*, vol. 3, no. 2, pp. 18–26.
- [5] Stoyneva, M. P., Uzunov, B. A., and Gärtner, G. (2015). Aerophytic green algae, epimycotic on *Fomes fomentarius* (L. ex Fr.) Kickx. *Annual of Sofia University "St. Kliment Ohridski", Faculty of Biology, Book 2*, vol. 99, pp. 19–25.

- [6] Neustroeva, N. V. and Mukhin, V. A. (2013). Symbiotic associations of xylotrophic basidiomycetes and algae, in *XIII Congress of Russian Botanical Society "Modern Botany in Russia" and Conference on Scientific Bases of Vegetation Protection and Rational Management in the Volga Basin*. Tolyatti: Cassandra.
- [7] Neustroeva, N. V., Kiseleva, I. S., and Mukhin, V. A. (2015). Carbon exchange between mycetobiont algae and wood-destroying fungi, in *All-Russian Conference with International Participation "Biodiversity and Ecology of Fungi and Mushroom-Like Organisms of Northern Eurasia"*. Ekaterinburg: Publishing House of the Ural University.
- [8] Mukhin, V. A., Patova, E. N., Kiseleva, I. S., et al. (2016). Mycetobiont symbiotic algae of wood-decomposing fungi. *Russian Journal of Ecology*, vol. 47, no. 2, pp. 133–137.
- [9] Mukhin, V. A., Patova, E. N., Sivkov M. D., et al. (2018). Diversity and Nitrogen-Fixing Activity of Phototrophic Mycetobionts of Xylotrophic Fungi. *Russian Journal of Ecology*, vol. 49, no. 5, pp.406–412.
- [10] Komárek, J. and Fott, B. (1983). *Chlorophyceae (Grünalgen). Ordnung: Chlorococcales, Das Phytoplankton des Süßwassers: Systematik und Biologie, 7*, part 1. Stuttgart: Das Phytoplankton des Süßwassers.
- [11] Ettl, H. and Gärtner, G. (2014). *Syllabus der Boden-, Luft- und Flechtenalgen*. Berlin & Heidelberg: Springer Spektrum.
- [12] Andreeva, V. M. (1998). *Soil and Aerophilic Green Algae (Chlorophyta: Tetrasporales, Chlorococcales, Chlorosarcinales)*. St. Petersburg: Nauka.
- [13] Neustroeva, N. V., Mukhin, V. A., Novakovskaya, I. V., et al. (2017). Host variability of mycetobiont algae. *Vestn. Udm. Univ., ser. "Biology. Earth Science"*, vol. 27, no. 3, pp. 291–296.
- [14] Neustroeva, N. V. and Mukhin, V. A. (2017). Symbiotic associations of xylotrophic fungi: Taxonomical and biomorphological composition mycetobiont algae, in *The Fourth Congress of Russian Mycologists: Current Mycology in Russia*. Moscow: Nat. Acad. Mycol.