

Content of ^{13}C and ^{15}N Isotopes in Bone Collagen of Geographical, Age, and Sex Groups of the Ural Cave Bear (Mammalia, Carnivora, Ursidae, *Ursus (Spelaearctos) kanivetzi* Verestchagin, 1973)

P. A. Kosintsev^{a,*}, K. Yu. Konovalova^a, and G. V. Simonova^b

Presented by Academician V.N. Bolshakov

Received August 8, 2023; revised August 30, 2023; accepted September 2, 2023

Abstract—Data on the content of ^{13}C and ^{15}N isotopes in the collagen of bones of the Ural cave bear (*Ursus (S.) kanivetzi* Verestchagin, 1973) from the North and Middle Urals were analyzed. The bones date from the first half of MIS 3. The bones of newborn individuals, individuals aged 1 year, males and females aged 2, 3, and 4 years, and older than 4 years were studied. Differences in $\delta^{13}\text{C}$ values between age, sex, and geographical samples are not significant. With age, the value of $\delta^{15}\text{N}$ significantly decreases, which is associated with weaning from milk nutrition to independent nutrition. The proportion of meat food in the diet of adult bears in the Middle Urals was higher than in the diet of adult bears in the North Urals. There are no noticeable differences in isotope signatures between males and females of different ages. The large cave bears of the Urals and Europe had a similar type of diet.

Keywords: *Ursus kanivetzi*, Ural cave bear, Late Pleistocene, the Urals, stable isotope, ^{13}C , ^{15}N , collagen

DOI: 10.1134/S001249662370076X

Analysis of the content of ^{13}C and ^{15}N isotopes in the tissues of living organisms is widely used to reconstruct the habitat and ecology of species in the past and present. It allows characteristics of the nutrition of the species and its position in the trophic chain to be evaluated. The study of isotopes is especially important for extinct species. One of the largest data sets on the isotopic composition of carbon and nitrogen in bone collagen was obtained for large cave bears (*Ursus (Spelaearctos) spelaeus* s.l.) of Western and Central Europe [1–7]. Data on the content of ^{13}C and ^{15}N isotopes in the collagen of the bones of large cave bears in Eastern Europe and the Urals are extremely scarce [8].

Analysis of morphological data and nuclear DNA showed that one species of large cave bear lived in Eastern Europe and the Urals in the late Pleistocene, i.e., the Ural cave bear (*Ursus (S.) kanivetzi* Verestchagin, 1973) [9, 10]. Three species lived in Central Europe: *U. (S.) spelaeus* Rosenmuller 1794, *U. (S.) eremus* Rabeder, Hofreiter, Nagel et Withalm, 2004, and

U. (S.) kanivetzi Verestchagin, 1973. In Western Europe, there were two species of large cave bears: *Ursus (S.) spelaeus* Rosenmuller 1794 and *U. (S.) eremus* Rabeder, Hofreiter, Nagel et Withalm, 2004 [10, 11].

The content of ^{13}C and ^{15}N isotopes in the collagen of the bones of the Ural cave bear from the Tain Cave in the Middle Urals [8] and the Medvezhiya Cave in the North Urals was studied. The selection from the Cave of Secrets has been supplemented with new specimens. Bear Cave (62°05' N, 58°05' E) is of karst origin, horizontal type of structure, length 480 m and height above sea level 280 m [12]. This is the northeastern edge of the range (*Ursus (S.) spelaeus* s.l.) [9]. More than 3000 bones of a large cave bear were found in the cave sediments, among which there are remains of all age groups from newborns to old individuals. This is a typical “graveyard” for cave bears, where the animals died during hibernation. Radiocarbon (AMS) dates were obtained from the cave bear bones: >48600 BP, no.? [13]; 42000 ± 450 BP, OxA-19608; 45150 ± 600 BP, OxA-19568 [10], which corresponds to the first half of Late Pleistocene marine isotope stage 3 (MIS 3). The Tain and Medvezhiya caves have the same taphonomic type (“cemetery”), the accumulation of bones occurred in one period—the first half of MIS 3 (57000–40000 years ago).

For analysis, 45 humeri, eight radii, and one tibia were taken. Determination of the sex and age of the

^aInstitute of Plant and Animal Ecology, Ural Branch, Russian Academy of Sciences, Yekaterinburg, Russia

^bInstitute of Monitoring of Climatic and Ecological Systems Siberian Branch Russian Academy of Sciences, Tomsk, Russia

*e-mail: kpa@ipae.uran.ru

Table 1. Values of δ¹³C and δ¹⁵N (‰) and standard deviation (sd) in collagen of the bones of the Ural cave bear (*U. (S.) kanivetz*) of the North and Middle Urals

Sex	Age	n	δ ¹³ C _{min}	δ ¹³ C _{max}	δ ¹³ C _{av} ± sd	δ ¹⁵ N _{min}	δ ¹⁵ N _{max}	δ ¹⁵ N _{av} ± sd
North Urals								
♂	>4+	9	-23.6	-21.2	-22.3±0.84	2.4	5.4	3.9±1.04
♀		5	-23.4	-21.8	-22.6±0.58	2.8	4.9	3.8±0.93
♂	3+, 4+	4	-22.2	-21.7	-21.9±0.21	3.9	5.8	4.8±0.84
♀		3	-23.6	-21.7	-22.4±1.01	3.4	4.2	3.9±0.44
♂	2+	4	-23.1	-21.9	-22.5±0.61	4.3	6.6	5.2±0.98
♀		4	-22.1	-20.2	-21.0±0.94	4.6	11.3	7.1±3.08
?	1+	23	-22.9	-21.4	-22.4±0.35	5.0	8.4	6.7±0.86
?	0+	2	-23.9	-22.4	-23.2±1.08	8.3	9.0	8.7±0.52
Middle Urals								
♂	>4+	5	-22.1	-21.3	-21.7±0.37	3.2	4.9	4.1±0.68
♀		2	-22.2	-21.8	-22.0±0.26	4.6	4.6	4.6
♂	3+, 4+	3	-22.5	-22.1	22.2±0.23	3.2	4.0	3.7±0.42
♀		4	-22.0	-21.4	-21.7±0.28	3.0	4.6	4.0±0.78
♂	2+	1			-22.8			7.2
♀		3	-22.1	-21.7	-21.8±0.23	5.4	5.7	5.6±0.15
?	1+	5	-22.8	-21.9	-22.5±0.34	4.8	7.8	6.4±1.11
?	0+	2	-24.2	-23.2	-23.7±0.71	7.5	9.0	8.3±1.06

¹ ♂—males, ♀—females, ?—sex not determined.

individuals to which the bones belonged was determined based on analysis of their size and the condition of the epiphyses (accreted or not accreted) [14–17]. Among them there are bones of males and females aged 2+, 3+, and 4+ years (subadults) and older than 4+ years (>4+, adults, adults) (Table 1). The sex of individuals aged 0+ and 1+ years is not determined. Each bone belongs to a separate individual. Individuals aged 0+ have only milk feeding, individuals aged 1+ have a mixed type of feeding (milk and independent), from age 2+ animals switch to independent feeding [18].

The isotopic composition of carbon (δ¹³C) and nitrogen (δ¹⁵N) in bone collagen was determined by isotope mass spectrometry using a DELTA V Advantage isotope mass spectrometer (Thermo Fisher Scientific, Germany) equipped with a Flash 2000 elemental analyzer (devices provided by the shared use center TomCKP SB RAS) according to standard methods. The VPDB standard has been adopted as the international carbon standard. Gaseous N₂ of atmospheric air is accepted as the international standard for nitrogen. Laboratory working gases for comparison, CO₂ and N₂, were calibrated according to the IAEA international standard sample—IAEA-600 Caffeine. The absolute measurement error of three consecutive measurements of the analyzed samples for δ¹³C did not exceed ±0.2‰, and for δ¹⁵N did not exceed ±0.4‰.

The values of the isotopic composition of carbon and nitrogen in the collagen of the bones of the Ural cave bear are presented in Table 1.

Mean δ¹³C values in groups of sub-adult and adult males and females in the North Urals differ by no more than 0.5‰, and δ¹⁵N values by no more than 0.9‰ (Table 1). In the Middle Urals, these differences for δ¹³C and δ¹⁵N are no more than 0.5‰ (Table 1). In general, the values of δ¹³C and δ¹⁵N for groups of sub-adult and adult individuals are close.

The differences in δ¹³C and δ¹⁵N values in groups of young individuals—2+, 1+, and 0+ years old—are much greater. The differences in δ¹³C values between males and females in the 2+ year group in the North Urals are 1.5‰, and in the Middle Urals – 1.0‰. The differences in the δ¹⁵N value are 1.9 and 1.6‰, respectively (Table 1). The differences in the δ¹³C value between groups 1+ and 0+ in the North Urals are 0.8‰, in the Middle Urals – 1.2‰, and the differences in the δ¹⁵N value are 2.0 and 1.9‰, respectively (Table 1).

The distribution of δ¹³C and δ¹⁵N values of age groups 1+ from the North and Middle Urals practically coincide and are almost completely outside the values for adult individuals (Fig. 1). The distribution of values for age groups 2+ of the North and Middle Urals coincide and partially overlap with the distribu-

tion of values for adult groups (Fig. 1). The values for age groups 2+ coincide with the values for age groups 1+. The values for individuals 0+ are outside the values for other age groups (Fig. 1).

Differences between samples (with $n > 4$) were statistically assessed using the nonparametric Mann-Whitney test. The $\delta^{13}\text{C}$ values do not differ statistically significantly between all samples. The $\delta^{15}\text{N}$ values differ significantly between the 4 pairs of samples. In the North Urals, $\delta^{15}\text{N}$ values are significantly (5% significance level) higher in the sample of age 1+ compared to the combined sample of males and females of age 2+ and in the combined sample of age 2+ compared to the combined sample of sub-adult (3+, 4+) males and females. In the Middle Urals, $\delta^{15}\text{N}$ values are significantly (1% significance level) higher in the sample of age 1+ than in the combined sample of subadult (+3, +4) and adult (>4+) males and females. In the combined sample of adult (>4+) males and females from the Middle Urals, the $\delta^{15}\text{N}$ values are significantly (5% significance level) higher than in a similar sample from the North Urals.

The data obtained show that almost all significant differences in $\delta^{15}\text{N}$ values are observed between samples of juveniles (1+ and 2+) or between samples of juveniles (2+) and subadults and adults (3+ and older). Differences in $\delta^{15}\text{N}$ values are determined by the proportion of protein foods in the diet [19]. High $\delta^{15}\text{N}$ values in younger age groups (0+, 1+) are associated with their dairy diet, which contains large amounts of protein. At the age of 2+, animals begin to feed independently, but the signature from the previous period of milk feeding is retained in the bone collagen. One individual aged 2+ has a very high $\delta^{15}\text{N}$ value—11.3 ‰. She may have continued to feed on milk into her second year of life.

There are reliably significant geographical differences between samples of adults. The $\delta^{15}\text{N}$ values for bears in the Middle Urals (4.5‰) are higher than those for bears in the North Urals (3.9‰) by 0.6‰. These differences indicate a difference in diet, in our case, a different ratio of plant and meat foods. In the Middle Urals, the share of meat food in the nutritional structure was higher than in the North Urals, but this difference is less than the differences between trophic levels [20].

The distribution of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in the collagen of the bones of the Ural cave bear almost completely coincides with the distribution of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in the collagen of the bones of cave bears (*U. (S.) spelaeus* and *U. (S.) eremus*) of Western and Central Europe [1–7] (Fig. 1). Outside the distribution there are three individuals, two of which are newborns (0+) and one young (2+) individual with an abnormally high $\delta^{15}\text{N}$ value. This indicates the similarity of the diet of all three species of cave bears, i.e., the Ural




Fig. 1. Distribution of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ (‰) values in the collagen of the bones of the Ural cave bear (*U. (S.) kanivetz*) of the North (blue) and Middle (red) Urals of different ages and large European cave bears (*U. (S.) spelaeus* s.l.).

(*U. (S.) kanivetz*) and the European (*U. (S.) spelaeus* and *U. (S.) eremus*).

Analysis of the data obtained shows the presence of significant age-related and geographical differences in the $\delta^{15}\text{N}$ values in the collagen of the bones of the Ural cave bear of the North and Middle Urals. There is a significant trophic shift between age groups associated with the transition from milk feeding to independent feeding. There are geographical differences in the diet of adults in the North and Middle Urals. The diet of bears in the Middle Urals contained a higher proportion of meat food. Differences in $\delta^{13}\text{C}$ values between age, gender and geographical samples are not significant. In general, adult and subadult males and females of the North and Middle Urals are at the same trophic level, because differences in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values (Table 1, Fig. 1) do not exceed the level of differences between different trophic levels [20]. The large cave bears of the Urals and Europe had a similar type of diet.

ACKNOWLEDGMENTS

We are grateful to the museum of the Institute of Plant and Animal Ecology of the Ural Branch of the Russian Academy of Sciences for providing samples for this study.

FUNDING

The study was supported by the Russian Science Foundation grant no. 22-24-01025, <https://rscf.ru/project/22-24-01025/>.

COMPLIANCE WITH ETHICAL STANDARDS

Conflict of interests. The authors declare that they have no conflicts of interest.

This article does not contain any studies involving animals performed by any of the authors.

OPEN ACCESS

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

REFERENCES

- Bocherens, H., Isotopic insights on cave bear palaeodiet, *Hist. Biol.*, 2019, vol. 31, no. 4, pp. 410–421.
- Robu, M., Fortin, J.K., Richards, M.P., et al., Isotopic evidence for dietary flexibility among European Late Pleistocene cave bears (*Ursus spelaeus*), *Can. J. Zool.*, 2013, vol. 91, no. 4, pp. 227–234.
- Bocherens, H., Stiller, M., Hobson, K.A., et al., Niche partitioning between two sympatric genetically distinct cave bears (*Ursus spelaeus* and *Ursus ingressus*) and brown bear (*Ursus arctos*) from Austria: Isotopic evidence from fossil bones, *Quat. Int.*, 2011, vol. 245, no. 2, pp. 238–248.
- Bon, C., Berthouaud, V., Fosse, P., et al., Low regional diversity of late cave bears mitochondrial DNA at the time of Chauvet Aurignacian paintings, *J. Archaeol. Sci.*, 2011, vol. 38, no. 8, pp. 1886–1895.
- Pérez-Rama, M., Fernández-Mosquera, D., and Grandal-d'Anglade, A., Recognizing growth patterns and maternal strategies in extinct species using stable isotopes: The case of the cave bear *Ursus spelaeus* ROSENMÜLLER, *Quat. Int.*, 2011, vol. 245, no. 2, pp. 302–306.
- Münzel, S.C., Stiller, M., Hofreiter, M., et al., Pleistocene bears in the Swabian Jura (Germany): Genetic replacement, ecological displacement, extinctions, and survival, *Quat. Int.*, 2011, vol. 245, no. 2, pp. 225–237.
- Nejman, L., Wood, R., Wright, D., et al., Hominid visitation of the Moravian Karst during the Middle-Upper Paleolithic transition: New results from Pod Hradem Cave (Czech Republic), *J. Hum. Evol.*, 2017, vol. 108, pp. 131–146.
- Kosintsev, P.A., Simonova, G.V., and Konovalova, K.Yu., First data on nutrition of the Ural cave bear *Ursus (Spelaearctos) kanivetz* Verestchagin, 1973 (Mammalia, Carnivora, Ursidae) as based on ¹³C and ¹⁵N isotope analyses, *Dokl. Biol. Sci.*, 2023, vol. 510, pp. 160–162.
- Gimranov, D.O. and Kosintsev, P.A., Cave bears (*Ursus spelaeus* sensu lato) of the Urals, *Paleontol. J.*, 2022, vol. 56, pp. 97–105.
- Rabeder, G., Hofreiter, M., Nagel, D., et al., New taxa of alpine cave bears (Ursidae, Carnivora), in *Cahiers Scientifiques-Museum d'Histoire Naturelle de Lyon*, 2004, vol. 2, pp. 49–68.
- Barlow, A., Pajmans, J.L.A., Federica, A., et al., Middle Pleistocene genome calibrates a revised evolutionary history of extinct cave bears, *Curr. Biol.*, 2021, vol. 31, no. 8, pp. 1771–1779.
- Guslitser, B.I. and Kanivets, V.I., *Peshchery pechorsko-go Urala* (Caves of the Pechora Urals), Moscow: Nauka, 1965.
- Baryshnikov, G.F., *Semeistvo medvezh'ikh (Carnivora, Ursidae)* (Family Ursidae (Carnivora)) St. Petersburg: Nauka, 2007.
- Vorob'ev, A.A., Dimensions of long tubular bones of cave bears of the Middle Urals, *Materialy konferentsii molodykh uchenykh "Sovremennye problemy populyatsionnoi, istoricheskoi i prikladnoi ekologii"* (Proc. Young Sci. Conf. "Current Problems in Population, Historical, and Applied Ecology"), Yekaterinburg: Ekaterinburg, 2001, pp. 38–41.
- Vorob'ev, A.A., Stages of postnatal ontogenesis of the skeleton of large cave bears, *Materialy konferentsii molodykh uchenykh "Biota gornykh territorii: Istoriya i sovremennoe sostoyanie"* (Proc. Young Sci. Conf. "Biota of Mountain Territories: History and Current State"), Yekaterinburg: Akademkniga, 2002, pp. 22–28.
- Kosintsev, P.A. and Vorob'ev, A.A., Biology of large cave bear (*Ursus spelaeus* Ros. et Hein.) in the Ural mountains, in *Mamont i ego okruzhenie: 200 let izucheniya* (The Mammoth and Its Ambience: 200 Years of Study), Yu.A., Rozanov, Ed., Moscow: Geos, 2001, pp. 266–278.
- Fosse, P. and Cregut-Bonnoure, E., Ontogeny/growth of (sub)modern brown bear (*Ursus arctos*) skeleton: A guideline to appraise seasonality for cave bear (*Ursus spelaeus*) sites?, *Quat. Int.*, 2014, vols. 339–340, pp. 275–288.
- Liden, K. and Angerbjorn, A., Dietary change and stable isotopes: A model of growth and dormancy in cave bears, *Proc. R. Soc. B*, 1999, vol. 266, pp. 1779–1783.
- Chisholm, B.S., Variation in diet reconstructions based on stable carbon isotopic evidence, in *The Chemistry of Prehistoric Human Bone*, Cambridge: Cambridge Univ. Press, 1989, pp. 10–37.
- Bocherens, H. and Drucker, D., Trophic level isotopic enrichment of carbon and nitrogen in bone collagen: Case studies from recent and ancient terrestrial ecosystems, *Int. J. Osteoarchaeol.*, 2003, vol. 13, nos. 1–2, pp. 46–53.

Translated by S. Nikolaeva

Publisher's Note. Pleiades Publishing remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.