

## Morphological Anomalies in *Rana arvalis* Nilsson, 1842 from Subarctic Urban Areas

V. L. Vershinin<sup>a, b, \*</sup>, S. D. Vershinina<sup>a</sup>, A. G. Trofimov<sup>a, b</sup>, and A. V. Burakova<sup>a</sup>

<sup>a</sup>Institute of Plant and Animal Ecology, Russian Academy of Sciences, Yekaterinburg, 620144 Russia

<sup>b</sup>Ural Federal University, Yekaterinburg, 620002 Russia

\*e-mail: vol\_de\_mar@list.ru

Received March 29, 2017

**Keywords:** *Rana arvalis*, polymorphism, morphological anomalies, the Subarctic, urbanization

**DOI:** 10.1134/S1067413618030116

The moor frog *Rana arvalis* Nilsson, 1842 is one of the most ecologically flexible, eurytopic species in Eurasia, with its range extending from the English Channel to Transbaikalia and from Kazakhstan to the Subarctic. In the Urals, this species is almost ubiquitous and occurs in both natural and anthropogenically transformed habitats [1–4]. The adaptive capacity of *R. arvalis* frogs living in the Far North “...reaches amazing perfection: even at about 0°C their digestive enzymes remain highly active, and the amplitude of heart contractions and the rate of ventricular tension development increase” [5]. Such an adaptive potential is largely related to specific features of polymorphism in this species [6–9].

Research on the population features of amphibians inhabiting the Subarctic was initiated in the 20th century [10], but deviant forms of morphogenesis and other qualitative indices characterizing specific features of population genetic structure have not yet been analyzed. The purpose of this study was to perform comparative analysis of the phenological (morphological) pattern of a new *R. arvalis* generation from urban areas of the Subarctic and Middle Urals.

The material was collected in the city of Labytnangi (the Polar Urals) in the zone of multistory buildings on Novaya St. on July 26–27, 2016. The total sample consisted of 69 postmetamorphic (juvenile) *R. arvalis* frogs, which were analyzed for qualitative and quantitative morphological parameters. Body length was measured with a digital caliper (Kraftool, Germany) to an accuracy of 0.01 mm; body weight, with a digital balance (Shimadzu, Japan) to an accuracy of 10 mg.

Two traits were used to analyze specific features of population genetic structure. The first was the frequency of the *striata* morph [11] phenotypically manifested as the presence of a light dorsomedial stripe extending to the tip of the snout. It is determined by the dominant allele of the diallelic autosomal gene

*striata* with complete dominance [12]. Specific features of inheritance make this trait a good marker of changes in the genetic structure of population. The second trait was the frequency of iris depigmentation, which is determined by a recessive mutation [13] whose expressivity and penetrance depend on the temperature during early stages of ontogeny [14].

Identification and analysis of exterior morphological anomalies were performed according to relevant classifications [15, 16] and also using original approaches [17]. The material for comparison was taken from the database for the territory of Yekaterinburg [18] in 1980 to 2013. Within this urban agglomeration, four zones with different habitat conditions for amphibians were distinguished [1]: (I) city center (not inhabited by amphibians) with multistory buildings, asphalt pavement, water bodies highly polluted by industrial emissions, and tubed small rivers and streams; (II) multistory districts with areas under development, wastelands, areas with bare soil, and highly polluted small water bodies; (III) low-rise districts of private houses with orchards and vegetable gardens, parks, and wasteland areas; (IV) park forest zone with multistory districts, areas under development, wastelands, areas with bare soil, and highly polluted small water bodies; habitats in this zone are mainly affected by recreational activities. The control sample (K) was from suburban populations living in a *Sphagnum*–horsetail pine forest 23 km from Yekaterinburg. The habitat in Labytnangi was also classified as a zone of multistory buildings. The acceptability of this typification is confirmed every year by hydrochemical analysis [18].

Statistical data processing was performed with Statistica for Windows 6.0.

Comparative morphological analysis of the sample from Labytnangi revealed no significant differences in body length, weight, and body condition from these parameters in juveniles from Yekaterinburg (Table 1),

**Table 1.** Morphological parameters of postmetamorphic *R. arvalis* frogs

Habitat (city)	Zone	<i>L</i> , mm	<i>P</i> , mg	<i>P/L</i> (body condition)	Number of animals
Labytnangi	II	16.4 ± 0.3	430.0 ± 22.6	25.2 ± 0.85	69
Yekaterinburg	II	16.4 ± 0.07	445.7 ± 4.9	26.7 ± 0.2	1561
Yekaterinburg	III	15.08 ± 0.09	345.1 ± 6.2	22.7 ± 0.3	999
Yekaterinburg	IV	15.02 ± 0.05	385.7 ± 3.7	22.5 ± 0.2	2844
Yekaterinburg	K	14.9 ± 0.07	368.8 ± 4.9	23.7 ± 0.2	1565

while the frequency of striata morph proved to be significantly higher than in juveniles from zone II in Yekaterinburg: 62.3 vs. 42.9% ( $\chi^2 = 12.88$ ;  $p = 0.0003$ ). The total frequency of animals with morphological anomalies was also significantly higher in Labytnangi than in the multistory zone of Yekaterinburg: 21.7 vs. 6.4% ( $\chi^2 = 25.65$ ;  $p \ll 0.0001$ ). The sample from Labytnangi was characterized by a high frequency of morphological anomalies (26.09%). In a decreasing order of occurrence, they include ectrodactyly, iris depigmentation, oligodactyly, abnormal skin pigmentation, and ectromelia (Table 2). For comparison: in the Middle Urals, the proportion of individuals with morphological anomalies among *R. arvalis* juveniles is 1.5–2% in forest populations versus 6–8% in urbanized areas. Iris depigmentation is a recessive mutation occurring with a frequency of about 0.39% (with a maximum of 3%) in forest populations and 1.7% (with a maximum of 7.5%) in urban populations; in the sample from Labytnangi, it was expressed bilaterally in 80% of mutants.

On the whole, a high total frequency of morphological anomalies was revealed among juveniles of new generation from *R. arvalis* population of the study area. Iris depigmentation accounts for a major proportion of all anomalies (33.3%), which is evidence for high homozygosity of this population. As shown for urban populations of amphibians [19–21], fragmentation of habitats and separation of populations into small isolates under the impact of urbanization lead to increasing homozygosity and higher rate of gene pool transformation due to genetic drift accompanied by inbreeding.

**Table 2.** Frequencies of morphological anomalies (%) in postmetamorphic *R. arvalis* from zones of multistory buildings

Parameter	City	
	Labytnangi	Yekaterinburg
Total number of animals	69	1561
Iris depigmentation	7.24	1.72
Abnormal skin pigmentation	2.9	0.94
Ectromelia	1.45	0.25
Ectrodactyly	11.6	0.25
Oligodactyly	2.9	0.16

The overall frequency of skeletal anomalies, which can be associated with trematode infestation [22], reaches 15.95%. However, only two species of intestinal parasites were found in juveniles of new generation from Labytnangi: nematodes *Oswaldocruzia filiformis* Goeze, 1782 and parasitic protists *Opalina ranarum* Dujardin, 1841. Skeletal anomalies observed in *R. arvalis* juveniles mainly included reduction of distal limb elements in different variants, probably of heterochrony type (Table 2). These results are preliminary and need verification in further studies.

A considerable proportion of striata morph revealed in *R. arvalis* population is evidence for the prevalence of individuals with genetically determined high reactivity of the hematopoietic system, several other physiological processes [6, 7, 9], rapid sexual maturation, and short life span [23]. It appears that the physiological specificity of this morph accounts for an increase in its frequency as an adaptive response of the population to anthropogenic transformation of the environment and to its geochemical background [24, 25] under conditions of strictly limited duration of the active life cycle phase in the Subarctic.

Thus, the results of this study show that *R. arvalis* population from the multistory zone of Labytnangi is characterized by a high level of specificity, which is probably due to insularization of the range and formation of isolates under the impact of urbanization combined with climatic features of the Subarctic.

#### ACKNOWLEDGMENTS

The study was performed within the frameworks of State Contract with the Institute of Plant and Animal Ecology, Ural Branch, Russian Academy of Sciences, and supported by the integrated program of the Ural Branch, Russian Academy of Sciences (project no. 18-4-4-28) and by Act 211 of the Government of the Russian Federation, contract no. 02.A03.21.0006.

#### REFERENCES

1. Vershinin, V.L., Distribution and species composition of amphibians within the city limits of Sverdlovsk, in *Informatsionnye materialy Instituta ekologii rastenii i zhivotnykh UrO AN SSSR* (Information Materials from Institute of Plant and Animal Ecology, Ural Branch, USSR Academy of Sciences), Sverdlovsk, 1980, pp. 5–6.

2. Vershinin, V.L., Preliminary assessment of the effect of anthropogenic factors on amphibians in Sverdlovsk, in *Problemy ekologii, ratsional'nogo ispol'zovaniya i okhrany prirodnykh resursov na Urale* (Problems of Ecology and Rational Management and Protection of Natural Resources in the Urals), Sverdlovsk, 1980, pp. 117–118.
3. Vershinin, V.L., Species composition and biological features of amphibians from several industrial cities in the Urals, *Extended Abstract of Cand. Sci. (Biol.) Dissertation*, Sverdlovsk, 1983.
4. Vershinin, V.L. and Toporkova, L.Ya., Amphibians in urban landscapes, in *Fauna Urala i Evropeiskogo Severa* (The Fauna of the Urals and Northern Europe), Sverdlovsk, 1981, pp. 48–56.
5. Shvarts, S.S., Evolutionary biology, *Vestn. Akad. Nauk SSSR*, 1974, no. 9, pp. 11–21.
6. Vershinin, V.L., Hematopoiesis in the Anura: Specific features of species adaptogenesis in recent ecosystems, *Entomol. Rev.*, 2004, vol. 84, Suppl. 1, pp. 113–119.
7. Vershinin, V.L., The striata morph in representatives of the genus *Rana* (Amphibia, Anura): Factors of adaptiveness to changes in the environment, *Zh. Obshch. Biol.*, 2008, vol. 69, no. 1, pp. 65–71.
8. Vershinin, V.L. and Vershinina, S.D., Comparative analysis of hemoglobin content in four species of anurans from the Ural Uplands, *Dokl. Biol. Sci.*, 2013, vol. 450, pp. 155–157.
9. Vershinin, V.L. and Vershinina, S.D., Physiological similarity of morphs due to homologous alleles in representatives of the Ranidae family, *Usp. Sovrem. Biol.*, 2013, vol. 113, no. 5, pp. 516–523.
10. Shvarts, S.S. and Ishchenko, V.G., *Puti prispobleniya nazemnykh pozvonochnykh zhitvnykh k usloviyam sushchestvovaniya v Subarktike* (Ways of Adaptation to Subarctic Conditions in Terrestrial Vertebrates), vol. 3: *Zemnovodnye* (Amphibians), *Tr. Inst. Ekol. Rast. Zhiv.*, vol. 79, Sverdlovsk: Ural. Fil. Akad. Nauk SSSR, 1971.
11. Schreiber, E., *Herpetologia europaea: Eine systematische Bearbeitung der Amphibien und Reptilien, welche bisher in Europa aufgefangen sind*, 2nd ed., Jena: Gustav Fisher, 1912.
12. Shchupak, E.L., Inheritance of dorsal band in *Rana arvalis*, *Informatsionnye materialy Instituta ekologii rastenii i zhitvnykh UrO AN SSSR* (Information Materials from Institute of Plant and Animal Ecology, Ural Branch, USSR Academy of Sciences), Sverdlovsk, 1977.
13. Dubois, A., Sur deux anomalies de la grenouille verte (*Rana esculenta*), *Bull. Soc. Linn. Lyon*, 1968, vol. 37, pp. 316–320.
14. Vershinin, V.L., Frequency of iris depigmentation in urban populations of *Rana arvalis* frogs, *Russ. J. Ecol.*, 2004, vol. 35, no. 1, pp. 58–62.
15. Dubois, A., Anomalies and mutations in natural populations of the *Rana "esculenta"* complex (Amphibia, Anura), *Mitt. Zool. Mus. Berlin*, 1979, no. 55, pp. 59–87.
16. Tyler, M.J., Natural pollution monitors, *Aust. Nat. Hist.*, 1983, vol. 21, pp. 31–33.
17. Vershinin, V.L., *Osnovy metodologii i metody issledovaniya anomalii i patologii amfibii: Ucheb. posobie* (Methodological Foundations and Methods of Studies on Anomalies and Pathologies of Amphibians: A Textbook), Yekaterinburg: Ural. Gos. Univ., 2015.
18. Vershinin, V.L., Vershinina, S.D., Berzin, D.L., et al., Long-term observation of amphibian populations inhabiting urban and forested areas in Yekaterinburg, Russia, *Sci. Data*, 2015, vol. 2, Article no. 150018, pp. 1–13. doi 10.1038/sdata.2015.18
19. Makeeva, V.M., Malyuchenko, O.P., Belokon', M.M., et al., Assessment of the state of gene pool in natural brown frog populations inhabiting human-transformed landscapes of the city of Moscow and Moscow region, in *Genetika v XXI veke: sovremennoe sostoyanie i perspektivy razvitiya* (Genetics in the 21st Century: Current state and Prospects for Development), Moscow, 2004, vol. 2, p. 211.
20. Makeeva, V.M., Belokon, M.M., Malyuchenko, O.P., and Leont'eva, O.A., Evaluation of the state of the gene pool of natural populations dwelling in the fragmented landscape of Moscow and Moscow district (with special reference to brown frogs), *Russ. J. Genet.*, 2006, vol. 42, no. 5, pp. 505–517.
21. Munshi-South, J., Zak, Y., and Pehek, E., Conservation genetics of extremely isolated urban populations of the northern dusky salamander (*Desmognathus fuscus*) in New York city, *PeerJ*, 2013, vol. 1, e64. doi 10.7717/peerj.64
22. Vershinin, V.L. and Neustroeva, N.S., The role of trematode infestation in the specifics of skeleton morphogenesis of *Rana arvalis* Nilsson, 1842 *Dokl. Biol. Sci.*, 2011, vol. 440, pp. 290–292.
23. Ledentsov, A.V., Dynamics of the age structure and size of the reproductive part of moor frog (*Rana arvalis* Nilss.) population, *Extended Abstract of Cand. Sci. (Biol.) Dissertation*, Sverdlovsk: Inst. Plant Anim. Ecol., Ural Branch, USSR Acad. Sci., 1990.
24. Sharygin, S.A., Trace elements in some amphibians and reptiles and their dynamics under the impact of anthropogenic factors, *Extended Abstract of Cand. Sci. (Biol.) Dissertation*, Sverdlovsk, 1980.
25. Vershinin, V.L., Specific features of the *Rana arvalis* Nilss. life cycle in the Eastern Ural Radioactive Trace, *Sib. Ekol. Zh.*, 2007, no. 4, pp. 677–682.

Translated by N. Gorgolyuk